

## WEAK IDEMPOTENT NIL-CLEAN RINGS

B. Asmare, T. Abebaw and K. Venkateswarlu\*

ABSTRACT. We introduce the concept of a weak idempotent nil-clean ring as a generalization of a weakly nil-clean ring. We give certain characterizations for weak idempotent nil-clean rings in terms of Jacobson radical and nil-radical. Further, we obtain any weak idempotent nil-clean ring  $R \cong R_1 \times R_2$  where  $R_1$  and  $R_2$  are weak idempotent nil-clean rings such that  $2 \in J(R_1)$  and  $3 \in J(R_2)$ .

### 1. INTRODUCTION

Throughout this paper,  $R$  stands for associative ring with unity unless and otherwise stated. We denote the set of all idempotents, nilpotents, units, the Jacobson radical, and the prime radicals (nil-radicals) of a ring  $R$  by  $Id(R)$ ,  $Nil(R)$ ,  $U(R)$ ,  $J(R)$  and  $N(R)$  respectively.

We recall the following definitions from [3]. A ring  $R$  is called

1. strongly nil-clean if for each  $r \in R$ , there exists a nilpotent  $n$  and an idempotent  $e$  such that  $r = n + e$  and  $ne = en$ .
2. nil-clean if every element can be expressed as a sum of a nilpotent and an idempotent.
3. strongly weakly nil-clean if each element  $r \in R$  can be represented as either  $r = n + e$  or  $r = n - e$ ,  $ne = en$  where  $n$  is nilpotent and  $e$  is idempotent.
4. weakly nil-clean ring if every element can be written as either a sum or a difference of a nilpotent and an idempotent.
5. clean if every element can be written as a sum of a unit and an idempotent.

The following hold: Strongly nil-clean  $\Rightarrow$  nil-clean  $\Rightarrow$  weakly nil-clean  $\Rightarrow$  clean.

It is observed that every element can be represented as a sum of a certain element and an idempotent element in all the above-said rings. It is quite natural to ask whether the representation can be generalized or not. In any

---

Published online: 1 April 2024

MSC(2020): Primary: 16N40; Secondary: 16N20, 16N99.

Keywords: Weakly nil-clean rings; Weak idempotent nil-clean rings; Clean rings.

Received: 27 May 2023, Accepted: 13 September 2023.

\*Corresponding author.

ring  $R$ , if  $a^4 = a^2$  then such  $a$  is called weak idempotent element. Clearly, every idempotent is weak idempotent but not conversely. For instance, consider the ring of integers modulo 4. Clearly, every element is a weak idempotent element but 2 is not idempotent. In view of these observations, is it possible to replace the idempotent element with a weak idempotent element in the above-said classes of rings? To some extent the answer is affirmative. In this context, we introduce the notion of weak idempotent nil-clean rings which is a subclass of the class of clean rings and a wider class to the class of weakly nil-clean rings.

In this paper, we introduce the notion of weak idempotent nil-clean rings (for short, win-clean rings) and furnish certain examples. Further, we obtain some basic results concerning weak idempotent nil-clean rings. In the next section, we prove  $R/Nil(R)$  is a reduced win-clean ring if and only if  $R$  is a commutative win-clean ring. Also, we characterize the win-clean ring in Proposition 2.23. The main result of this paper is that every win-clean ring  $R$  is isomorphic to a direct product of win-clean rings  $R_1$  and  $R_2$  where  $2 \in J(R_1)$  and  $3 \in J(R_2)$ .

## 2. MAIN RESULTS

**Definition 2.1.** Let  $R$  be a ring. An element  $a \in R$  is called weak idempotent nil-clean if  $a = n + w$  for some nilpotent  $n$  and some weak idempotent  $w$ .  $R$  is said to be weak idempotent nil-clean if every element of  $R$  is weak idempotent nil-clean.

*Remark 2.2.* We denote the set of all weak idempotent elements by  $wi(R)$  and weak idempotent nil-clean ring by win-clean ring.

**Example 2.3.** Let  $R = M_2(\mathbb{Z}_3)$ . Then  $R$  is win-clean ring.

**Example 2.4.** Let  $R = \mathbb{Z}_3 \times \mathbb{Z}_3$ . Then  $R$  is win-clean ring.

*Remark 2.5.* If  $R$  is a ring and  $w$  is a weak idempotent element, then

- (1)  $w^{2n} = w^2$ , and  $w^{2n+1} = w^3$ .
- (2)  $Id(R) \cup -Id(R) \subseteq wi(R)$ .

We can easily verify that every weakly nil-clean ring is a win-clean ring using remark 2.5 (2) but the converse is not true. For instance,  $\mathbb{Z}_3 \times \mathbb{Z}_3$  is win-clean ring but not weakly nil-clean, since  $(2, 1)$  cannot be expressed as a sum or a difference of any nilpotent and any idempotent element in  $\mathbb{Z}_3 \times \mathbb{Z}_3$ .

**Theorem 2.6.** *Let  $R$  be a ring. If  $w \in R$  is weak idempotent, then*

- (1)  $w^k$  is a weak idempotent element, i.e,  $w^{2k} = w^{4k}$  where  $k \in \mathbb{N}$ .
- (2)  $w^2$  and  $1 - w^2$  are idempotent elements.
- (3)  $2w^2 - 1$  and  $w - 1 + w^2$  are units.
- (4)  $w^n - w^{n+2}$  is nilpotent for every  $n \in \mathbb{N}$ .
- (5)  $(1 - w^2)w^2 = 0$
- (6)  $w$  is clean.

*Proof.* It is straightforward. □

**Definition 2.7.** Let  $R$  be a ring and  $I$  be an ideal of  $R$ . Then the weak idempotents can be lifted modulo  $I$  if there exists  $w \in wi(R)$  for a given  $a \in R$  with  $a^4 - a^2 \in I$  such that  $w - a \in I$ .

**Proposition 2.8.** Let  $I$  be a nil ideal of a ring  $R$ . If  $\bar{w}$  is a weak idempotent element in  $R/I$ , then  $\bar{w}$  can be lifted to a weak idempotent in  $R$ .

*Proof.* Let  $\bar{w} \in R/I$  be a weak idempotent and  $w$  be any pre-image for  $\bar{w}$ . Then  $\bar{w}^2 = \bar{w}^4$  implies that  $w^2 - w^4 \in I$  or  $w^2 \equiv w^4 \pmod{I}$  where  $w^2$  and  $w^4$  are pre-images of  $\bar{w}^2$  and  $\bar{w}^4$ , in  $R/I$  respectively. Let  $z = 1 - w^2$ . Then (a)  $w^2z = zw^2$  and (b)  $w^2 + z \equiv 1 \pmod{I}$ .

Now  $w^2z = w^2 - w^4 \in I$ . Then  $0 = (w^2z)^k = w^{2k}z^k$  for some positive integer  $k$ . Also,  $w^{2k}$  is a pre-image of  $\bar{w}$ , since  $w^{2k} \equiv w^2 \pmod{I}$ . Conditions (a) and (b) are preserved when  $w$  and  $z$  are replaced by  $w^{2k}$  and  $z^k$ . Moreover, condition (c)  $w^2z = zw^2 = 0$  is also preserved.

From condition (b), we have  $x = 1 - w^2 - z \in I$ . Then  $(1 - w^2 - z)^m = 0$  for some positive integer  $m$ . Thus  $1 = 1 - x^m = (1 - x)(1 + x + \cdots + x^{m-1})$  and it follows that  $1 - x$  has an inverse  $u = 1 + x + \cdots + x^{m-1}$ .  $u$  commutes with  $w$  and  $z$  as  $x$  commutes with  $w$  and  $z$ .

Since  $x \in I$ ,  $u \equiv 1 \pmod{I}$ . We can replace  $w$  and  $z$  with  $uw^2$  and  $uz$ , in this case  $w$  is again a pre-image for  $\bar{w}$  and also conditions (a), (b), and (c) hold true. Further, it is true that (d)  $w^2 + z = 1$ . By condition (c), we have  $w^2z = 0$ , so it gives that  $w^2 = w^2(w^2 + z) = w^4 + w^2z = w^4$ . Therefore,  $\bar{w}$  lifted to the weak idempotent  $w$  in  $R$ . □

**Proposition 2.9.** The homomorphic image of any win-clean ring is win-clean.

*Proof.* It is straightforward. □

*Remark 2.10.* The converse of Theorem 2.9 is not true. For instance, consider the canonical epimorphism  $\alpha : \mathbb{Z} \rightarrow \mathbb{Z}/(3)$  given by  $\alpha(n) = n + (3)$ . Then  $\mathbb{Z}_3 \cong \mathbb{Z}/(3)$  is a win-clean ring, but  $\alpha^{-1}(\mathbb{Z}/(3)) = \mathbb{Z}$  is not a win-clean ring.

Let  $R$  be a ring and  $M$  a left  $R$ -module. Consider the idealization of  $R$  and  $M$  given by  $R(M) = R \oplus M$ . For  $(r, m), (s, t) \in R(M)$ , product and sum defined as follows:

$$(r, m)(s, t) = (rs, rt + sm); (r, m) + (s, t) = (r + s, m + t).$$

Then  $R(M)$  is the ring.

**Theorem 2.11.** *Let  $R$  be a ring and  $M$  be a left  $R$ -module. Then  $R$  is win-clean if and only if  $R(M)$  is win-clean.*

*Proof.* Assume that  $R$  is win-clean ring and  $(r, m) \in R(M)$  where  $r \in R$  and  $m \in M$ . Then  $r = n + w$  for  $n \in Nil(R)$  and  $w \in wi(R)$ . Thus  $n^k = 0$  for  $k \in \mathbb{N}$ . So  $(n, m)^{k+1} = (n^{k+1}, (k+1)n^k m) = (0, 0)$  which implies that  $(r, m) = (n + w, m) = (n, m) + (w, 0)$  is win-clean expression of  $(r, m)$ . Hence,  $R(M)$  is win-clean. Conversely,  $R \cong R(M)/(0 \oplus M)$  is homomorphic image of  $R(M)$ . So by Theorem 2.9,  $R$  is win-clean ring.  $\square$

**Proposition 2.12.** *Let  $R$  be a ring. Then weak idempotent elements in  $J(R)$  are nilpotents.*

*Proof.* Let  $w \in J(R)$  be a weak idempotent element. Then  $w^2 \in J(R)$  and also  $1 - w^2$  is an idempotent element. Again,  $w^2 \in J(R)$  implies that  $1 - w^2 \in U(R)$ . So  $1 - w^2$  is both idempotent and unit. Thus  $1 - w^2 = 1$ , since 1 is the only unit and idempotent element. This implies that  $w^2 = 0$ . Hence  $w$  is nilpotent element.  $\square$

**Proposition 2.13** ([8]). *Let  $R$  be a ring and  $a, b \in R$  such that  $ab \neq ba$ . Then*

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k} + \sum_{k=0}^n D_k b^{n-k}$$

where  $d_a(x) = ax - xa$  and  $D_k = D_k(b, a) = (a + d_b)^n 1 - a^n$ ,  $D_0(b, a) = 0$ ,  $D_{n+1}(b, a) = d_b a^n + (A + d_b) D_n(b, a)$ .

**Proposition 2.14.** *Let  $R$  be a win-clean ring, then  $J(R) \subseteq Nil(R)$ .*

*Proof.* Let  $a \in J(R)$ . Then  $a = n + w$ , where  $n \in Nil(R)$  and  $w \in wi(R)$ . Then  $(a - w)^k = 0$  for some  $k \in \mathbb{N}$ . So  $(w - a)^k \in J(R)$ . Now

$$(w - a)^k = \sum_{k=0}^n \binom{n}{k} w^k a^{n-k} + \sum_{k=0}^n D_k a^{n-k}$$

implies that  $(w - a)^k - [\sum_{k=0}^{n-1} \binom{n}{k} w^k a^{n-k} + \sum_{k=0}^n D_k a^{n-k}] = w^k \in J(R) \cap wi(R)$ . Since  $J(R)$  does not contain units and non-zero idempotents,  $w$  must be nilpotent. Now  $a - w, w \in Nil(R)$  which in turn implies that  $a \in Nil(R)$ . Hence  $J(R) \subseteq Nil(R)$ .  $\square$

**Corollary 2.15.** *If  $R$  is a win-clean ring, then  $J(R)$  is nil.*

*Remark 2.16.* A reduced win-clean ring is a ring in which all the elements are weak idempotents.

**Proposition 2.17.** *Let  $R$  be a commutative ring. Then  $R$  is win-clean if and only if  $R/Nil(R)$  is a reduced win-clean ring.*

*Proof.* Assume that  $R$  is win-clean ring. Let  $\bar{x} = x + Nil(R) \in R/Nil(R)$  for some  $x \in R$ . Now write

$$\bar{x} = (n + w) + Nil(R) = (n + Nil(R)) + (w + Nil(R)) = w + Nil(R)$$

and  $w + Nil(R) \in wi(R/Nil(R))$ . This implies that  $\bar{x}$  is weak idempotent element in  $R/Nil(R)$ . Since  $\bar{x}$  is arbitrary,  $R/Nil(R)$  is reduced win-clean ring. Conversely, assume that  $R/Nil(R)$  is win-clean ring and let  $r \in R$ . Since  $R/Nil(R)$  is reduced,  $Nil(R/Nil(R)) = \{0\}$  and  $r + Nil(R) = w + Nil(R)$  for some  $w + Nil(R) \in wi(R/Nil(R))$ . Then  $w^4 - w^2 \in Nil(R)$ . By Proposition 2.12, the weak idempotent  $w + Nil(R)$  can be lifted to a weak idempotent  $w \in wi(R)$  such that  $r - w = n$  for some  $n \in Nil(R)$ , i.e.,  $r = n + w$ . This shows that  $r$  is win-clean. Hence  $R$  is win-clean ring.  $\square$

**Corollary 2.18.** *Let  $R$  be a commutative ring. Then  $R$  is win-clean if and only if  $R/N(R)$  is win-clean ring.*

*Proof.* It is obvious.  $\square$

**Proposition 2.19.** *Let  $I$  be a nil ideal of a ring  $R$ .  $R$  is win-clean if and only if  $R/I$  is win-clean.*

*Proof.* ( $\implies$ ) It is obvious.

( $\impliedby$ ) Let  $r \in R$ . Then  $\bar{r} = r + I \in R/I$ . We can write  $\bar{r} = \bar{n} + \bar{w}$  where  $\bar{n} \in Nil(R/I)$  and  $\bar{w} \in wi(R/I)$  implies that  $r + I = (n + w) + I$ . The nilpotent  $\bar{n}$  in  $R/I$  lift to a nilpotent  $n$  in  $R$ . To see this,  $\bar{n}^k = 0$  for  $k \geq 1$  in  $R/I$  implies that  $n^k \in I$ . Since  $I$  is nil,  $(n^k)^m = 0$ . So  $n^{km} = 0$  for  $m \geq 1$ . We know that weak idempotents lift modulo any nil ideal, this allows us to assume that  $w$  is a weak idempotent in  $R$ . Moreover,  $r - n - w \in I$ . It follows that  $r - w = n + d$  where  $d \in I$ . Since  $n^m = 0$  for some  $m \in \mathbb{N}$ , we have  $(n + d)^k \in I$  because  $I$  is ideal of  $R$ . Thus  $(n + d)^{mk} = 0$  for some  $m \in \mathbb{N}$  as  $I$  is nil ideal. So  $n + d$  is nilpotent. Therefore,  $R$  is win-clean, as desired.  $\square$

**Corollary 2.20.** *A ring  $R$  is win-clean if and only if  $R/J(R)$  is win-clean and  $J(R)$  is nil.*

*Proof.* Since  $J(R)$  is nil, the proof follows from Proposition 2.19.  $\square$

The converse of Proposition 2.14 is not true. Consider example 1.2 in [7]. If we take a simple domain  $F = \mathbb{Z}_5$ , then  $A = M_2(\mathbb{Z}_5)$  is a ring of  $2 \times 2$  matrices over integer modulo 5, and  $B = D_2(\mathbb{Z}_5)$  is a ring of  $2 \times 2$  diagonal matrices over integer modulo 5 such that  $Nil(B) = \begin{pmatrix} 0 & \mathbb{Z}_5 \\ 0 & 0 \end{pmatrix}$ . Define  $R = B + A[[x]]_x$ , where  $A[[x]]$  denotes the formal power series ring with an indeterminate  $x$  over a ring  $R$ . Then  $Nil(R) \subsetneq J(R) = Nil(B) + A[[x]]$  and  $R/J(R) \cong \mathbb{Z}_5$ . But  $\mathbb{Z}_5$  is not win-clean and hence  $R/J(R)$  is not win-clean. Therefore, By Corollary 2.20,  $R$  is not win-clean ring.

*Remark 2.21.* It is clear that if  $x \in R$  a non-zero central nilpotent, then  $1 - xr \in U(R)$  for all  $r \in R$ . Hence  $x \in J(R)$ , i.e, the non-zero central nilpotents are contained in Jacobson radical,  $J(R)$ .

**Corollary 2.22.** *Let  $R$  be a win-clean ring such that the weak idempotents are central. Then  $C(R)$ , the center of  $R$ , is a win-clean ring.*

**Proposition 2.23.** *The following are equivalent for a ring  $R$ :*

- (1)  $R$  is win-clean.
- (2)  $12$  is nilpotent and  $R/12R$  is win-clean.
- (3)  $R/J(R)$  is win-clean and  $J(R)$  is nil.

*Proof.* (1)  $\implies$  (2). If  $12 = 0$ , then we are done. Assume that  $12 \neq 0$ . As  $R$  is a ring with 1,  $1 + 1 = 2 \in R$  is the least non-unit central element of  $R$ . Then there exist a weak idempotent  $w$  and a nilpotent  $n$  such that  $2 = n + w$ . Thus  $(2 - n)^2 = (2 - n)^4 \implies 2^2 - 4n + n^2 = 2^4 - 32n + 24n^2 - 8n^3 + n^4$ . So  $n(-n^3 + 8n^2 - 23n + 28) = 12$ . Hence,  $12$  is nilpotent. Since  $R$  is win-clean,  $R/12R$  is win-clean by Proposition 2.19.

(2)  $\implies$  (1) follows from Proposition 2.19 and (1)  $\iff$  (3) obtained immediately from Corollary 2.20.  $\square$

**Proposition 2.24** ([5]). *Let  $R$  be a ring, and let  $I$  be any nil-ideal of  $R$ . Then  $R$  is nil-clean if and only if  $R/I$  is nil-clean.*

**Proposition 2.25.** *A ring  $R$  is nil-clean if and only if  $R$  is win-clean and  $2 \in J(R)$ .*

*Proof.* ( $\implies$ ) Suppose  $R$  is nil-clean and  $r \in R$ . Then  $r = n + e$  where  $n \in Nil(R)$  and  $e \in Id(R)$ . Thus  $e \in wi(R)$ . So  $r$  is win-clean and hence  $R$  is win-clean. Also,  $2 = n + e$  implies that  $n = 2$ . Thus  $2$  is central nilpotent. This implies that  $2 \in J(R)$ .

( $\Leftarrow$ ) Assume  $R$  is win-clean. Then  $J(R)$  is nil. As  $2 \in J(R)$ ,

$$2 + J(R) = 0 + J(R).$$

We know that a nilpotent modulo nil ideal lifted to nilpotent in  $R$ . So we have  $2 = 0$ , i.e.,  $\text{char}(R/J(R)) = 2$ . Thus for all  $r \in R$  we have  $2\bar{r} = \bar{0}$  and  $1 - 2\bar{r} = \bar{1}$ . So  $R/J(R)$  is Boolean. Hence  $R/J(R)$  is nil-clean. By Proposition 2.24,  $R$  is nil-clean.  $\square$

**Proposition 2.26.** *A ring  $R$  is weakly nil-clean if and only if  $R$  is win-clean and  $2 \in J(R)$  or  $3 \in J(R)$ .*

*Proof.* ( $\Rightarrow$ ) Obviously  $R$  is win-clean. Assume that  $2 \notin J(R)$ . Then  $2 \in U(R)$  and  $6^n = 0$  for some positive integer  $n$  as 6 is nilpotent element in  $R$  ([3], Theorem 2). Thus  $2^n 3^n = 0$  implies that  $3^n = 0$ . Hence  $3 \in J(R)$ .

( $\Leftarrow$ ) Assume that  $R$  is win-clean. Then  $R/J(R)$  is win-clean and  $J(R)$  is nil by Corollary 2.20. If  $2 \in J(R)$ , then by Proposition 2.25,  $R$  is nil-clean. So  $R$  is weakly nil-clean. Again, if  $3 \in J(R)$ , then  $3 + J(R) = 0 + J(R)$  and also 2 is invertible in  $R$ . we can assume  $3 = 0$ , so that  $\text{char}(R/J(R)) = 3$ . So  $\bar{2}$  is unit in  $R/J(R)$ . Moreover,  $3\bar{r} = \bar{0}$ ,  $\bar{1} - 3\bar{r} = \bar{1}$  and  $\bar{2} - 3\bar{r} = \bar{2}$  for all  $r \in R$ . Thus  $R/J(R) \cong \mathbb{Z}_3$  and hence  $R/J(R)$  is weakly nil-clean. Therefore,  $R$  is weakly nil-clean.  $\square$

**Proposition 2.27.** *A finite direct product  $R = \prod R_\alpha$  of rings is win-clean ring if and only if each  $R_\alpha$  is win-clean ring.*

*Proof.* It is straightforward.  $\square$

**Proposition 2.28.** *Let  $R$  be a ring. Then  $R$  is win-clean ring if and only if  $R \cong R_1 \times R_2$  where  $R_1$  is win-clean with  $2 \in J(R_1)$  and  $R_2$  is 0 or a win-clean ring with  $3 \in J(R_2)$ .*

*Proof.* ( $\Rightarrow$ ) Suppose  $R$  is win-clean ring. Then 12 is nilpotent element in  $R$ , so that  $(12)^n = 0$  for some positive integer  $n$ . Then  $4^n R \cap 3^n R = 0$  and  $4^n R + 3^n R = R$ . Thus  $R \cong (R/2^{2n}R) \times (R/3^n R)$  by Chinese remainder theorem. By Proposition 2.27,  $R_1 = R/2^{2n}R$  and  $R_2 = R/3^n R$  are win-clean rings. Thus 2 is central nilpotent in  $R_1$ . So  $2 \in J(R_1)$ . We can assume  $R_2 \neq 0$ . Then 3 is central nilpotent in  $R_2$  and hence  $3 \in J(R_2)$ .

( $\Leftarrow$ ) It is obvious.  $\square$

**Corollary 2.29.** *The following are equivalent for a ring  $R$ .*

- (1)  $R$  is a win-clean ring with central weak idempotent elements.

- (2)  $R \cong R_1 \times R_2$ , where  $R_1$  is win-clean with weak idempotents are central and  $J(R_1)$  nil such that  $R_1/J(R_1)$  is Boolean, and  $R_2$  is 0 or  $R_2/J(R_2) \cong \mathbb{Z}_3$  with  $J(R_2)$  nil.
- (3)  $R$  is win-clean ring with central weak idempotent elements,  $J(R)$  is nil, and  $R/J(R)$  is isomorphic to either a Boolean ring, or to  $\mathbb{Z}_3$ , or to the direct product of two such rings.

*Proof.* (1)  $\implies$  (2) Using Proposition 2.28, we can write  $R \cong R_1 \times R_2$ , where  $R_1$  is win-clean ring with central weak idempotents and  $2 \in J(R_1)$ ; and  $R_2$  is 0 or win-clean ring with central weak idempotents and  $3 \in J(R_2)$ . Thus  $\text{char}(R_1/J(R_1)) = 2$  which in turn implies that  $\bar{x} = -\bar{x}$  for all  $\bar{x} \in R_1/J(R_1)$ . Hence,  $R_1/J(R_1)$  is Boolean. Assume  $R_2 \neq 0$ . As  $R_2$  is win-clean and  $3 \in J(R_2)$ ,  $R_2/J(R_2)$  is win-clean and  $\text{char}(R_2/J(R_2)) = 3$ . Also, 2 is unit in  $R_2$ , since  $2 \notin J(R_2)$ . From this, we conclude that

$$R_2/J(R_2) = \{3R_2, 1 - 3R_2, 2 - 3R_2\},$$

so that every element of  $R_2/J(R_2)$  is nilpotent or invertible. Therefore,  $R_2/J(R_2) \cong \mathbb{Z}_3$ . Furthermore, by Corollary 2.20,  $J(R_1)$  and  $J(R_2)$  are nil ideals.

(2)  $\implies$  (3) and (3)  $\implies$  (1) are straightforward.  $\square$

**Theorem 2.30.** *Let  $R$  be a reduced commutative ring. The following statements are equivalent.*

- (1)  $R = \text{wi}(R)$ .
- (2)  $R$  is isomorphic to either a Boolean ring  $B$  or  $\mathbb{Z}_3$ , or  $B \times \mathbb{Z}_3$ .
- (3) For all  $x \in R$ ,  $x^4 = x^2$ .
- (4)  $R$  is win-clean.

*Proof.* For a reduced ring  $R$ , (1)  $\iff$  (3)  $\iff$  (4). Thus, it remains to show the equivalence of (1) and (2).

(1)  $\implies$  (2) Suppose  $R = \text{wi}(R)$ . If  $y \in R$ ,  $y^2$  is an idempotent. If  $R$  is indecomposable, then either  $y^2 = 0$  or  $y^2 = 1$  for any  $y \in R$ . This implies that  $y = 0$  or  $y^2 = 1$  for all  $y \in R$ . Thus, each nonzero element of  $R$  is a unit and hence  $R$  is a field. Hence,  $R$  is isomorphic to either  $\mathbb{Z}_2$  or  $\mathbb{Z}_3$ .

Next, assume  $R$  is not indecomposable. Let  $R = S \times T$  and  $s \in S$ , where  $S$  and  $T$  are coprime ideals of  $R$ , that is,  $S + T = R$ . Then,  $(s, 0)$  is not a unit implies that either  $(s, 0) = (0, 0)$ , or  $(s, 0)^2 = (0, 0)$ , or  $(s, 0)^2 = (s, 0)$ , or  $(s, 0)^2 = (s, 0)^4$  and  $(s, 0)^2 \neq (1, 0)$ . If  $(s, 0) = (0, 0)$  or  $(s, 0)^2 = (0, 0)$ , then  $(s, 0) = (0, 0)$  since  $S$  is reduced. In this case,  $S$  is a field. So,  $S$  is isomorphic to  $\mathbb{Z}_2$  or  $\mathbb{Z}_3$ . If  $(s, 0)^2 = (s, 0)$ , then  $s \in \text{Id}(S) \cup [-\text{Id}(S)]$  and



hence  $S = Id(S) \cup [-Id(S)]$ . By [4, Theorem 1.13],  $S$  is isomorphic to either a Boolean ring, or  $\mathbb{Z}_3$ , or  $B \times \mathbb{Z}_3$ , where  $B$  is a Boolean ring. The same holds for  $T$ . As a direct product of two Boolean rings is a Boolean ring we get  $R$  is isomorphic to a Boolean ring  $B$ ,  $\mathbb{Z}_3 \times \mathbb{Z}_3$ ,  $\mathbb{Z}_3 \times B$ , or  $\mathbb{Z}_3 \times \mathbb{Z}_3 \times B$ . If  $(s, 0)^2 = (s, 0)^4$  and  $(s, 0)^2 \neq (1, 0)$ , then let  $y = (s, 0)$ . Now  $R = R(y^2) \oplus R(1 - y^2)$  is the decomposition of  $R$ .

Assume  $R(y^2)$  is not a Boolean ring. Then, we show that  $R(1 - y^2)$  is Boolean. Suppose  $ry^2$  is not idempotent. Then, for any  $s \in R$ ,  $ry^2 + (-s)(1 - y^2)$  is not idempotent. Thus,  $-(ry^2 + (-s)(1 - y^2)) = -ry^2 + s(1 - y^2)$  is idempotent. So, each  $s(1 - y^2)$  is idempotent. Thus,  $R(1 - y^2)$  is Boolean and also  $2R(1 - y^2) = 0$ . Hence, for each  $y \in R$ , either  $2y^2 = 0$  or  $2(1 - y^2) = 0$ .

If  $(0 : 2) = \{y \in R \mid 2y^2 = 0\} = R$ , then  $char(R) = 2$ . Hence,  $R = wi(R) = Id(R)$  and so  $R$  is Boolean. Now assume  $(0 : 2) \neq R$ . Then, we claim that  $(0 : 2)$  is a maximal ideal of  $R$ . Suppose there is a maximal ideal  $M$  such that  $(0 : 2) \subseteq M$ . Let  $y^2 \in M - (0 : 2)$ . Then,  $y^2 \in wi(R) = R$  and  $y^2 \notin (0 : 2)$ . Thus,  $2y^2 \neq 0$  and hence  $2(1 - y^2) = 0$ . So,  $1 - y^2 \in (0 : 2) \subseteq M$ , a contradiction. Hence,  $(0 : 2)$  is a maximal ideal. So,  $\bar{R} = R/(0 : 2)$  is an indecomposable ring with  $\bar{R} = wi(\bar{R})$ . By the idea in the first part of this proof, we have that  $\bar{R}$  is isomorphic to  $\mathbb{Z}_2$  or  $\mathbb{Z}_3$ .

Next we show that  $2R \cap (0 : 2) = 0$ . Assume that  $y \in 2R \cap (0 : 2)$ . Then,  $y = 2s$  and  $2y^2 = 0$ . But then  $y^2 = y^4 = (2s)^4 = 2(2s)^2(2s)^2 = 2y^2 = 0$ . If  $2R = 0$ , then  $R$  is Boolean.

Now assume that  $2R \neq 0$ . If  $2R = R$ , then  $(0 : 2) = 0$  is a maximal ideal of  $R$ . Thus,  $R$  is a field and hence by the first paragraph of this proof, it is isomorphic to  $\mathbb{Z}_3$ . If  $2R \neq R$ , then  $R = 2R \oplus (0 : 2)$ , where  $(0 : 2)$  is a Boolean ring and  $2R \cong R/(0 : 2)$  is isomorphic to  $\mathbb{Z}_3$  since  $2R \cong \mathbb{Z}_2$  or  $\mathbb{Z}_3$  by the first paragraph of this proof and  $2R \not\subseteq (0 : 2)$ . Therefore,  $R$  is isomorphic to either a Boolean ring, or  $\mathbb{Z}_3$ , or  $B \times \mathbb{Z}_3$ , or  $\mathbb{Z}_3 \times \mathbb{Z}_3$ , or  $\mathbb{Z}_3 \times \mathbb{Z}_3 \times B$ , where  $B$  is a Boolean ring.

(2)  $\implies$  (1) It is obvious. □

Recall that a ring is said to be zero dimensional if every prime ideal is maximal ideal.

**Corollary 2.31.** *Let  $R$  be a commutative ring. The following statements hold.*

- (1) *A reduced indecomposable ring is win-clean if and only if it is isomorphic to either  $\mathbb{Z}_2$  or  $\mathbb{Z}_3$ . In particular, any win-clean domain is isomorphic to either  $\mathbb{Z}_2$  or  $\mathbb{Z}_3$ .*

(2) *A win-clean ring is zero-dimensional.*

*Proof.* (1) Suppose  $R$  is a reduced indecomposable win-clean. Then 0 is the only nilpotent, and its idempotents are only 0 and 1. Let  $w \in wi(R)$ . Then  $w^2 \in Id(R)$  implies that  $w^2 = 0$  or  $w^2 = 1$ . If  $w^2 = 0$ , then  $w$  is both weak idempotent and nilpotent. So  $w = 0$ . If  $w^2 = 1$ , then  $w$  is a unit and weak idempotent. Now we have  $R = \{0, 1, w\}$ . Since  $R$  is closed under  $+$ ,  $w + 1 \in R$  which implies that  $w + 1 = 0$  or  $w + 1 = 1$ , or  $w + 1 = w$ . If  $w + 1 = 0$ , then  $w = -1$ . In this case,  $R = \{0, 1, -1\}$  which is isomorphic to  $\mathbb{Z}_3$ . If  $w + 1 = 1$  or  $w + 1 = w$ , then  $w = 0$  as  $0 \neq 1$ . Hence  $R = \{0, 1\}$  which is isomorphic to  $\mathbb{Z}_2$ . The converse is straightforward.

(2) Let  $R$  be a win-clean and  $P$  a prime ideal of  $R$ . Then  $R/P$  is an integral domain. By (1), the quotient  $R/P$  is isomorphic to either  $\mathbb{Z}_2$  or  $\mathbb{Z}_3$  and hence  $P$  is maximal ideal. □

**Proposition 2.32.** *Let  $R$  be a win-clean ring with central weak idempotent elements and let  $a \in R$ . If  $aR$  contains no non-zero idempotent. Then  $a$  is the sum of two nilpotent elements.*

*Proof.* Suppose  $aR$  contains no non-zero idempotent. Choose  $w \in wi(R)$  and  $n \in Nil(R)$  such that  $a = n + w$ . Then

$$aw^3 = nw^3 + w^4 = nw^3 + w^2 = (nw + 1)w^2.$$

So  $aw^3(nw + 1)^{-1} = (nw + 1)w^2(nw + 1)^{-1}$  in  $aR$ . As  $nw$  is nilpotent,  $nw + 1$  is unit and  $w^2$  is idempotent. Thus  $(nw + 1)w^2(nw + 1)^{-1}$  is idempotent. Since  $aR$  does not contain non-zero idempotent element, we have

$$(nw + 1)w^2(nw + 1)^{-1} = 0$$

which implies  $w^2 = 0$  and hence  $w$  is nilpotent. Therefore,  $a$  is a sum of two nilpotent elements. □

**Definition 2.33.** Let  $R$  be a ring. Then an element  $x$  in  $R$  is called the square root of idempotent element if there exists an idempotent element  $e$  in  $R$  such that  $x^2 = e$ .

**Proposition 2.34.** *Let  $R$  be a win-clean ring with central weak idempotent elements in which,  $2 \in U(R)$ . Then every element of  $R$  can be written as a sum of nilpotent and a square root of idempotent element.*

*Proof.* Let  $a \in R$ . Then  $a = n + w$  for some  $n \in Nil(R)$  and  $w \in wi(R)$ . Let  $v = 2w^2 - 1$ . Then  $v^2 = (2w^2 - 1)^2 = 4w^4 - 4w^2 + 1 = 4w^2 - 4w^2 + 1 = 1$ . Thus  $vv^{-1} = (2w^2 - 1)(2w^2 - 1)^{-1} = 1$ . Now  $v = 2w^2 - 1$  implies  $w^2 = (v + 1)/2$  and  $[(v + 1)/2]^2 = (v + 1)/2$ . Therefore,  $w$  is a square root of idempotent.  $\square$

Next, we see that a win-clean ring is a subclass of clean rings.

**Theorem 2.35.** *Every win-clean ring is clean.*

*Proof.* Let  $R$  be a win-clean ring and  $a \in R$ . Then  $a = n + w$  for some nilpotent  $n$  and weak idempotent  $w$ . So  $a = n + w = (n + w - 1 + w^2) + (1 - w^2)$ . By Theorem 2.6,  $w - 1 + w^2$  is unit and  $1 - w^2 \in Id(R)$ . To see  $n + w - 1 + w^2$  is unit. Let  $u = w - 1 + w^2$ . Then  $n + w - 1 + w^2 = n + u$ . Since  $n$  and  $(u^{-1}n)$  are nilpotents, we have  $n^m = 0$  and  $(u^{-1}n)^m = 0$  for some positive integer  $m$ . Now

$$\begin{aligned} (n + u)^{-1} &= [u(1 + \frac{n}{u})]^{-1} = [1 - \frac{n}{u} + (\frac{n}{u})^2 - (\frac{n}{u})^3 + \dots + (-\frac{n}{u})^{m-1}]u^{-1} \\ &= [1 - u^{-1}n + (u^{-1}n)^2 - \dots + (-1)^{m-1}(u^{-1}n)^{m-1}]u^{-1}. \end{aligned}$$

and so

$$\begin{aligned} (n + u)(n + u)^{-1} &= (n + u)[1 - u^{-1}n + \dots + (-1)^{m-1}(u^{-1}n)^{m-1}]u^{-1} \\ &= n[1 - u^{-1}n + \dots + (-1)^{m-1}(u^{-1}n)^{m-1}]u^{-1} \\ &\quad + u[1 - u^{-1}n + (u^{-1}n)^2 - \dots + (-1)^{m-1}(u^{-1}n)^{m-1}]u^{-1} \\ &= nu^{-1} - (nu^{-1})^2 + (nu^{-1})^3 - \dots + (-1)^{m-1}(nu^{-1})^m \\ &\quad + 1 - nu^{-1} + (nu^{-1})^2 - \dots + (-1)^{m-1}(nu^{-1})^{m-1} \\ &= 1. \end{aligned}$$

Thus  $a$  is clean. Therefore,  $R$  is clean.  $\square$

In general, the converse of Theorem 2.35 does not hold true. For example, integer modulo 5,  $\mathbb{Z}_5$ , is clean but not win-clean.

**Lemma 2.36.** *If  $w$  is a weak idempotent element in a win-clean ring  $R$  and  $2 \in J(R)$ , then  $w \pm w^2$  is nilpotent.*

*Proof.* Since  $2 \in J(R)$ , we have  $(w \pm w^2)^2 = 2(w^2 \pm w^3) \in J(R)$ . As  $J(R)$  is nil, there exists some positive integer  $m$  such that  $2^m = 0$  and also  $(w \pm w^2)^{2m} = 0$ . Hence,  $w \pm w^2$  is nilpotent.  $\square$

The following proposition sets a condition for which a clean element becomes win-clean.

**Proposition 2.37.** *Let  $R$  be a commutative ring,  $2 \in J(R)$  and  $x$  be clean in  $R$  with clean decomposition  $x = u + e$ . Then  $x$  is win-clean if and only if there exists  $w \in wi(R) \cap Nil(R)$  such that  $2e - 1 + u$  is nilpotent.*

*Proof.* ( $\implies$ ) Suppose  $x$  is win-clean. Then  $x = n + f$  for some  $n \in Nil(R)$  and  $f \in wi(R)$ . Now  $x = n + f = (n - 1 + f + f^2) + (1 - f^2)$ . Since  $2 \in J(R)$ ,  $f + f^2$  is nilpotent by Lemma 2.36. Then take  $u = n - 1 + f + f^2$  and  $e = 1 - f^2$ . So

$$2e - 1 + u = 2(1 - f^2) - 1 + (n - 1 + f + f^2) + f^2 = n + f.$$

( $\impliedby$ ) We can rewrite  $x = u + e$  as  $x = (u + 2e - 1 + w^2) + (1 - e - w^2)$ . Since  $1 - e - w^2$  is weak idempotent,  $x$  is win-clean.  $\square$

### Acknowledgments

We thank the unknown reviewers for their invaluable suggestions and comments for the improvement of this paper.

### REFERENCES

1. S. Ali, A note on commutative weakly nil clean rings, *J. Algebra Appl.*, **15**(10) (2016), Article ID: 1620001.
2. D. K. Basnet and J. Bhattacharyya, Weak nil-clean rings, (2015), <https://doi.org/10.48550/arXiv.1510.07440>.
3. S. Breaz, P. Danchev and Y. Zhou, Rings in which every element is either a sum or a difference of a nilpotent and an idempotent, *J. Algebra Appl.*, **15**(08) (2016), Article ID: 1650148.
4. P. V. Danchev and W. W. McGovern, Commutative weakly nil clean unital rings, *J. Algebra*, **425** (2015), 410–422.
5. A. J. Diesl, Nil clean rings, *J. Algebra*, **383** (2013), 197–211.
6. T. Koşan, Z. Wang, and Y. Zhou, Nil-clean and strongly nil-clean rings, *J. Pure Appl. Algebra*, **220**(2) (2016), 633–646.
7. C. I. Lee and S. Y. Park, When nilpotents are contained in Jacobson radicals, *J. Korean Math. Soc.*, **55**(5)(2018), 1193–1205.
8. W. Wyss, Two non-commutative binomial theorems, (2017), <https://doi.org/10.48550/arXiv.1707.03861>
9. M. I. Zubayda and N. F. Norihan, On weak nil-clean rings, *Open Access Library Journal*, **9**(e8812) (2022), 2333–9721.

**Biadiglign Asmare**

Department of Mathematics, College of Natural and Computational Science, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia.

Email: [biadiglign.asmare@aau.edu.et](mailto:biadiglign.asmare@aau.edu.et)

**Tilahun Abebaw**

Department of Mathematics, College of Natural and Computational Science, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia.

Email: [tilahun.abebaw@aau.edu.et](mailto:tilahun.abebaw@aau.edu.et)

**Kolluru Venkateswarlu**

Department of Computer Science and System Engineering, College of Engineering, Andhra University, Visakhapatnam, Andhra Pradesh, India.

Email: [drkvenkateswarlu@gmail.com](mailto:drkvenkateswarlu@gmail.com)

WEAK IDEMPOTENT NIL-CLEAN RINGS

B. ASMARE, T. ABEBAW AND K. VENKATESWARLU

حلقه‌های پوچ-تمیز خودتوان ضعیف

بی. اسماره<sup>۱</sup>، تی. آباوا<sup>۲</sup> و کی. ونکتسوارلو<sup>۳</sup>

<sup>۱,۲</sup>گروه ریاضیات، کالج علوم طبیعی و محاسباتی، دانشگاه آدیس آبابا، آدیس آبابا، اتیوپی

<sup>۳</sup>گروه علوم کامپیوتر و مهندسی سیستم، کالج مهندسی، دانشگاه آندرا، ویساخاپاتنام، آندرا پرادش، هند

در این مقاله، مفهوم حلقه پوچ-تمیز خودتوان ضعیف را که تعمیمی از حلقه‌های پوچ-تمیز ضعیف است، تعریف می‌کنیم. با توجه به جیکوبسون رادیکال و پوچ-رادیکال، یک مشخصه سازی خاص برای حلقه‌های پوچ-تمیز خودتوان‌های ضعیف ارائه می‌دهیم. به علاوه، نشان می‌دهیم که برای هر حلقه پوچ-تمیز خودتوان ضعیف  $R$  داریم  $R \cong R_1 \times R_2$  که  $R_1$  و  $R_2$  حلقه‌های پوچ-تمیز خودتوان ضعیف هستند به طوری که  $2 \in J(R_1)$  و  $3 \in J(R_2)$ .

کلمات کلیدی: حلقه‌های پوچ-تمیز ضعیف، حلقه‌های پوچ-تمیز خودتوان ضعیف، حلقه‌های تمیز.